# Extreme values of the Dedekind $\Psi$ function

Patrick Solé Michel Planat

arXiv:1011.1825v2 [math.NT] 18 Jan 2011

**Abstract**

Let $\Psi(n) := n\prod_{p|n}(1+\frac{1}{p})$ denote the Dedekind $\Psi$ function. Define, for $n \geq 3$, the ratio $R(n) := \frac{\Psi(n)}{n\log\log n}$. We prove unconditionally that $R(n) < e^{\gamma}$ for $n \geq 31$. Let $N_n = 2\cdots p_n$ be the primorial of order $n$. We prove that the statement $R(N_n) > \frac{e^{\gamma}}{\zeta(2)}$ for $n \geq 3$ is equivalent to the Riemann Hypothesis.

**MsC codes:** 11N56, 11A25, 11M06

**Keywords:** Dedekind $\Psi$ function, Euler totient function, Mertens formula, Nicolas bound, Primorial numbers

## I. Introduction

The Dedekind $\Psi$ function is an arithmetic multiplicative function defined for every integer $n > 0$ by

$$\Psi(n) := n\prod_{p|n}(1+\frac{1}{p}).$$

It occurs naturally in questions pertaining to dimension of spaces of modular forms [2] and to the commutation of operators in quantum physics [6]. It is related to the sum of divisor function

$$\sigma(n) = \sum_{d|n} d$$

by the inequalities

$$\Psi(n) \leq \sigma(n),$$

and the fact that they coincide for $n$ squarefree. It is also related to Euler $\varphi$ function by the inequalities

$$n^2 > \varphi(n)\Psi(n) > \frac{n^2}{\zeta(2)}$$

derived in Proposition 5 below.

In view of the studies of large values of $\sigma$ [7] and of low values of $\varphi$ [4], it is natural to study both the large and low values of $\Psi$. To that end, we define the ratio $R(n) := \frac{\Psi(n)}{n\log\log n}$. The motivation for this strange quantity is the asymptotics of Proposition 3. We prove unconditionally that $R(n) < e^{\gamma}$, for $n \geq 31$ in Corollary 2. Note that this bound would follow also from the Robin inequality

$$\sigma(n) \leq e^{\gamma} n\log\log n$$

for $n \geq 5041$ under Riemann Hypothesis (RH) [7], since $\Psi(n) \leq \sigma(n)$.
In the direction of lower bounds, we prove that the statement $R(N_n) > \frac{e^{\gamma}}{\zeta(2)}$ for $n \geq 3$ is equivalent to RH, where $N_n = 2\cdots p_n$ is the primorial of order $n$. The proof relies on Nicolas's work on the Euler totient function [4].

Telecom ParisTech, 46 rue Barrault, 75634 Paris Cedex 13, France.
Institut FEMTO-ST, CNRS, 32 Avenue de l'Observatoire, F-25044 Besançon, France

## II. Reduction to primorial numbers

Define the primorial number $N_n$ of index $n$ as the product of the first $n$ primes

$$N_n = \prod_{k=1}^{n} p_k,$$

so that $N_1 = 2,\ N_2 = 6, \cdots$ and so on. As in [4], the primorial numbers play the role here of superabundant numbers in [7]. They are champion numbers (ie left to right maxima) of the function $x \mapsto \Psi(x)/x$ :

$$\frac{\Psi(m)}{m} < \frac{\Psi(n)}{n} \text{ for any } m < n, \tag{1}$$

We give a proof of this fact, which was observed in [5].

*Proposition 1:* The primorial numbers $N_n$ are exactly the champion numbers of the function $x \mapsto \Psi(x)/x$.

*Proof:* The proof is by induction on $n$. The induction hypothesis $H_n$ is that the statement is true up to $N_n$. It is clear that $H_2$ is true. Let $N_n \leq m < N_{n+1}$ be a generic integer. The number $m$ has at most $n$ distinct prime factors. This, in combination with the observation that $1 + 1/x$ is monotonically decreasing as a function of $x$, shows that $\Psi(m)/m \leq \Psi(N_n)/N_n$. Further $\Psi(N_n)/N_n < \Psi(N_{n+1})/N_{n+1}$. The proof of $H_{n+1}$ follows. ■

In this section we reduce the maximization of $R(n)$ over all integers $n$ to the maximization over primorials.

*Proposition 2:* Let $n$ be an integer $\geq 2$. For any $m$ in the range $N_n \leq m < N_{n+1}$ one has $R(m) \leq R(N_n)$.

*Proof:* Like in the preceding proof we have

$$\Psi(m)/m \leq \Psi(N_n)/N_n$$

Since $0 < \log\log 6 < \log\log N_n \leq \log\log m$, the result follows. ■

## III. $\Psi$ at primorial numbers

We begin with an easy application of Mertens formula [3, Th. 429].

*Proposition 3:* We have, as $n \to \infty$

$$\lim R(N_n) = \frac{e^\gamma}{\zeta(2)} \approx 1.08.$$

*Proof:* Writing $1 + 1/p = (1 - 1/p^2)/(1 - 1/p)$ in the definition of $\Psi(n)$ we can combine the Eulerian product for $\zeta(2)$ with Mertens formula

$$\prod_{p \leq x} (1 - 1/p)^{-1} \sim e^\gamma \log(x)$$

to obtain

$$\frac{\Psi(N_n)}{N_n} \sim \frac{e^\gamma}{\zeta(2)} \log(p_n),$$

Now the Prime Number Theorem [3, Th. 6, Th. 420] states that $x \sim \theta(x)$ for $x$ large. where $\theta(x)$ stands for Chebyshev's first summatory function:

$$\theta(x) = \sum_{p \leq x} \log p.$$

This shows that, taking $x = p_n$ we have

$$p_n \sim \theta(p_n) = \log(N_n).$$

The result follows. ∎

This motivates the search for explicit upper bounds on $R(N_n)$ of the form $\frac{e^\gamma}{\zeta(2)}(1+o(1))$. In that direction we have the following bound.

*Proposition 4:* For $n$ large enough to have $p_n \geq 20000$, that is $n \geq 2263$, we have

$$\frac{\Psi(N_n)}{N_n} \leq \frac{\exp(\gamma + 2/p_n)}{\zeta(2)}(\log\log N_n + \frac{1.125}{\log p_n})$$

So, armed with this bound, we derive a bound of the form $R(N_n) < e^\gamma$ for $n \geq A$, with $A$ a constant.

*Corollary 1:* For $n \geq 4$, we have $R(N_n) < e^\gamma = 1.78\cdots$

*Proof:*

For $p_n \geq 20000$, we use the preceding proposition. We need to check that

$$\exp(2/p_n)(1 + \frac{1.125}{\log(p_n)\log\log(N_n)}) \leq \zeta(2).$$

Since the LHS is a decreasing function of $n$ it is enough to check this inequality for the first $n$ such that $p_n \geq 20000$.

For $5 \leq p_n \leq 20000$, that is $3 \leq n \leq 2262$ we simply compute $R(N_n)$, and check that it is $< e^\gamma$. ∎

We can extend this Corollary to all integers $> 30$ by using the reduction of preceding section, combined with some numerical calculations for $30 < n \leq N_4$.

*Corollary 2:* For $n > 30$, we have $R(n) < e^\gamma$.

We prepare for the proof of the preceding Proposition by a pair of Lemmas. First an upper bound on a partial Eulerian product from [8, (3.30) p.70].

*Lemma 1:* For $x \geq 2$, we have

$$\prod_{p \leq x}(1 - 1/p)^{-1} \leq e^\gamma(\log x + \frac{1}{\log x})$$

Next an upper bound on the tail of the Eulerian product for $\zeta(2)$.

*Lemma 2:* For $n \geq 2$ we have

$$\prod_{p > p_n}(1 - 1/p^2)^{-1} \leq \exp(2/p_n)$$

*Proof:* Use Lemma 6.4 in [1] with $x = p_n$ and $t = 2$. ∎

We are now ready for the proof of Proposition 4.

*Proof:*

Write

$$\frac{\Psi(N_n)}{N_n} = \prod_{k=1}^{n} \frac{1 - 1/{p_k}^2}{1 - 1/p_k}$$

and use both lemmas to derive

$$\frac{\Psi(N_n)}{N_n} \leq \frac{\exp(\gamma + 2/p_n)}{\zeta(2)}(\log p_n + \frac{1}{\log p_n}).$$

Now we get rid of the first $\log$ in the RHS by the bound of [7, p.206]

$$\log(p_n) < \log\log N_n + \frac{0.125}{\log p_n}.$$

The result follows. ∎

## IV. Lower bounds

We reduce first to Euler's $\varphi$ function.

*Proposition 5:* For $n \geq 2$ we have

$$n^2 > \varphi(n)\Psi(n) > \frac{n^2}{\zeta(2)}$$

*Proof:* The first inequality follows at once upon writing

$$\frac{\varphi(n)\Psi(n)}{n^2} = \prod_{p|n}(1 - 1/p^2),$$

a product of finitely many terms $< 1$. Notice for the second inequality that

$$\frac{\varphi(n)\Psi(n)}{n^2} = \prod_{p|n}(1 - 1/p^2) > \prod_{p}(1 - 1/p^2),$$

an infinite product that is the inverse of the Eulerian product for $\zeta(2)$. ∎

*Theorem 1:* Under RH the ratio $R(N_n)$ is $> \frac{e^\gamma}{\zeta(2)}$ for $n \geq 3$. If RH is false, this is still true for infinitely many $n$.

*Proof:*

Follows by Proposition 5, combined with [4, Theorem 2]. ∎

In view of this result and of numerical experiments the natural conjecture is

*Conjecture 1:* For all $n \geq 3$ we have $R(N_n) > \frac{e^\gamma}{\zeta(2)}$.

The main result of this note is the following.

*Theorem 2:* Conjecture 1 is equivalent to RH.

*Proof:* If RH is true we refer to the first statement of Theorem 1. If RH is false we consider the function

$$g(x) := \frac{e^\gamma}{\zeta(2)} \log\theta(x) \prod_{p\leq x}(1 + 1/p)^{-1},$$

Observing that $\log\theta(p_n) = \log\log N_n$, we see that $g(p_n) < 1$ is equivalent to $R(N_n) > \frac{e^\gamma}{\zeta(2)}$. We need to show that there exists an $x_0 \geq 3$ such that $g(x_0) > 1$ or equivalently $\log g(x_0) > 0$. Using once again the identity $1 - 1/p^2 = (1 - 1/p)(1 + 1/p)$, and [1, Lemma 6.4], we obtain, upon writing

$$-\log\zeta(2) = \sum_{p\leq x}\log(1 - 1/p^2) + \sum_{p>x}\log(1 - 1/p^2),$$

the bound

$$\log g(x) \geq \log f(x) - 2/x,$$

where $f$ is the function introduced in [4, Theorem 3], that is

$$f(x) := e^\gamma \log\theta(x) \prod_{p\leq x}(1 - 1/p).$$

We know by [4, Theorem 3 (c)] that, if RH is false, there is a $0 < b < 1$ such that $\limsup x^{-b} f(x) > 0$ and hence $\limsup \log f(x) >> \log x$. Since $2/x = o(\log x)$, the result follows. ∎

## V. Conclusion

In this note we have derived upper and lower bounds on the Dedekind $\Psi$ function. We show unconditionally that the function $\Psi(n)$ satisfies the Robin inequality. Since $\psi(n) \leq \sigma(n)$ this could be proved under $RH$ [7] or by referring to [1]. Of special interest is Conjecture 1 which is shown here to be equivalent to RH. We hope this new RH criterion will stimulate research on the Dedekind $\Psi$ function.

**Acknowledgements:** The authors thank Fabio Anselmi, Pieter Moree, and Jean-Louis Nicolas for helpful discussions.

## References

[1] Choie, YoungJu, Lichiardopol, Nicolas, Moree, Pieter, Solé, Patrick On Robin's criterion for the Riemann hypothesis, J. Théor. Nombres Bordeaux 19 (2007), no. 2, 357–372.

[2] J. A. Csirik, M. Zieve and J. Wetherell, On the genera of $X_0(N)$, unpublished manuscript (2001); available online at `http://www.csirik.net/papers.html`

[3] G.H. Hardy, E.M. Wright, *An introduction to the theory of numbers*, Oxford (1979).

[4] Nicolas, Jean-Louis Petites valeurs de la fonction d'Euler. J. Number Theory 17 (1983), no. 3, 375–388.

[5] Michel Planat, Riemann hypothesis from the Dedekind psi function, `hal.archives-ouvertes.fr/docs/00/52/64/54/PDF/RiemannHyp.pdf`

[6] Michel Planat, Pauli graphs when the Hilbert space dimension contains a square: why the Dedekind psi function ? J. Phys. A: Math. Theor. 44 (2011) 045301-16.

[7] G. Robin, Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann. J. Math. Pures Appl. (9) 63 (1984), 187–213.

[8] J. B. Rosser and L. Schoenfeld, Approximate formulas for some functions of prime numbers. Illinois J. Math. 6 (1962), 64–94.